Recognition and distribution of *Solanum hoplopetalum* and *S. hystrix* (Solanaceae) in Australia

Solanum hoplopetalum Bitter & Summerh. and S. hystrix R. Br. are two closely related species of Solanum subgenus Leptostemonum. The first named taxon, Solanum hystrix, was described by Brown (1810) from plants he collected in South Australia, to which the species is confined. Early collections of a related taxon endemic to Western Australia were initially also treated as S. hystrix (e.g. Black 1926), prior to their recognition as a distinct species (S. hoplopetalum) by Bitter & Summerhayes (1926). However, Bitter & Summerhayes' new taxon was apparently overlooked or ignored by later workers (e.g. Grieve & Blackall 1975, Meadly 1960 and Parsons 1967), who continued to treat Western Australian populations as S. hystrix. A recent revision of Solanum in Australia (Symon 1981) clarified the status of these taxa, but it is apparent that, at least in Western Australia, some degree of confusion still exists as to the recognition of the two species, and on a wider scale, their occurrence outside their respective 'natural' ranges.

Both taxa are also among the very few indigenous solanums that could be considered (minor) agricultural weeds (Parsons & Cuthbertson 1992, Symon 1981), so accurate identification of the species involved is important.

Recognition of the species

The main character separating *S. hoplopetalum* and *S. hystrix* is indumentum of the mature vegetative parts, this best observed on the abaxial leaf surface. Hairs are also present on reproductive structures in both taxa, but are not so useful for identification. *S. hoplopetalum* and *S. hystrix* are unusual in that their indumentum is composed of apparently simple hairs, as most members of subgenus *Leptostemonum* have an indumentum comprised of characteristic stellate hairs. The simple hairs in *S. hoplopetalum*, *S. hystrix* and some other unrelated taxa of subgenus *Leptostemonum* are probably derived from stellate hairs, where the lateral rays have been gradually reduced to the point of being 'lost' altogether, or are very poorly developed (i.e. reduced-stellate hairs); the phylogenetic significance of these hairs is discussed in Lepschi & Symon (in press).

Leaf dimensions and distribution also aid in identification, but are not exclusive. The two taxa may be distinguished by the following key:



Figure 1. A-B Solanum hystrix. A - indumentum on abaxial leaf surface (x10), B - simple (reduced stellate) hairs (left), and simple glandular hairs (right) (x330). C-D Solanum hoplopetalum. C - indumentum on abaxial leaf surface (x10), D - simple (reduced stellate) hairs (left), and simple glandular hairs (right) (x330). Drawn from Lepschi & Lally 1735 (A-B) and Keighery 4363 (C-D).

Species treatments

Recent morphological descriptions and notes on the nomenclature of both taxa can be found in Purdie *et. al* (1982) and Symon (1981), and a description of *S. hystrix* is also in Symon (1986). For information on the weed biology of *S. hoplopetalum*, see Meadly (1960) and Paterson (1967), and for both taxa Parsons & Cuthbertson (1992).

Solanum hoplopetalum Bitter & Summerh.

Distribution. Occurs mainly in the drier parts of south-western Western Australia, from at least Wiluna, south to Lake Grace and Newdegate, east to Queen Victoria Springs and Caiguna, and westwards to Coorow, Toodyay and Wagin, with sporadic introduced occurrences in the more humid regions of the far south-west corner, e.g. Perth metropolitan area (G.J. Keighery pers. comm.) and Ludlow. Meadly (1960) reports the species from Geraldton and Borden, but no specimens have been seen. This species has almost certainly benefitted from agricultural expansion in Western Australia during this century (*cf.* Bitter & Summerhayes 1926), and it is not known whether some populations at the extremes of the distribution represent natural occurrences or more recent introductions.

An isolated introduced population has been recorded from South Australia, but has not persisted.

Selected specimens examined. Natural distribution: WESTERN AUSTRALIA: Cundelee, 1967, *P. Boswell* R11 (PERTH); Tammin sand plain, 2/1953, *B.J. Grieve s.n.* (PERTH); 8.5 km E of Caiguna Roadhouse, 17/2/1995, *B.J. Lepschi & T.R. Lally* 1741 (AD, CANB, PERTH); 4 km N of Koolyanobbing on road to Helena & Aurora Ranges, 29/9/1995, *B.J. Lepschi* 2084 (CANB, NSW, PERTH); 10 km SE of Ardath, 28/9/1977, *R.D Pearce* 119 (AD).

Adventive occurrences. SOUTH AUSTRALIA: Coombe Siding, 1/1951, *leg. ign.* and Coombe Station, 2/1940, *E.S. Alcock s.n.* (AD mixed sheet); Coombe Siding, 7/2/1941, *E.S. Alcock s.n.* (AD). WESTERN AUSTRALIA: Ludlow, 27/12/1994, *G.J. Keighery* 13341 (PERTH).

Habitat. S. hoplopetalum occurs most frequently in disturbed sites such as roadsides, railway lines and grazing land, although it also inhabits open woodland, mallee and scrub communities, frequently on heavier soils (e.g. loam, clay-loam) rather than pure sands. As with many other *Solanum* species, it responds well to fire, and it can be common in recently burnt areas.

Solanum hystrix R. Br.

Distribution. Eyre Peninsula region of South Australia, extending from the eastern margin of the Nullarbor Plain, about 40 km west-north-west of Yalata Roadhouse, east to near Iron Knob, north to at least the Transcontinental Railway Line and south to the coast (including offshore islands). Like *S. hoplopetalum, S. hystrix* may have benefited from agriculture and land clearance. Recorded once from south-eastern South Australia, New South Wales and Western Australia, but not persisting.

Selected specimens examined. Natural distribution: SOUTH AUSTRALIA: South Australian Railway Line, 1 mile [c. 2 km] N of Cummins, 19/3/1969, C.R. Alcock 2660 (AD, CANB, G, H, IBSC, MA); St Peters Island, Nuyts Archipelago, 19/10/1988, L. Jansen NPIS965 (AD); 64.5 km NE of Kimba on road to Iron Knob, 15/2/1995, B.J. Lepschi & T.R. Lally 1735 (AD, CANB, NSW, PERTH); 8.5 km WNW of Yalata Roadhouse on road to Nullarbor Roadhouse, 16/2/1995, B.J. Lepschi & T.R. Lally 1737 (AD, CANB); Gawler Range, NW of Minnipa in vicinity of Scrubby Peak Homestead, 4/10/1972, D.E. Symon 8366 (AD).

Adventive occurrences: SOUTH AUSTRALIA: 5 km S of Port Wakefield, 4/12/1981, R. Majury s.n. (AD 98570418). WESTERN AUSTRALIA: 100 miles [c. 160 km] E of Norseman, 20/12/1968, G. Howard s.n. (AD, CANB, PERTH).

Habitat. Similar to that described for S. hoplopetalum, although S. hystrix also occurs in coastal (littoral) sites and sandy soils (Symon 1981).

Occurrence as introduced plants

Neither species is known to be established as an alien outside of its respective state. *S. hoplopetalum* appears to have persisted at Coombe in South Australia for eleven years (1940-1951), unless there were repeated introductions which seems unlikely. Searches by D.E. Symon at this site in recent years have failed to relocate it (see also Symon 1986).

The single records of *S. hystrix* from northern New South Wales (Conn 1992; Symon 1981), southeastern South Australia (Purdie *et al.* 1982) and Western Australia (Symon 1981) all appear to have been chance introductions, none of which appears to have established a permanent population. Of course, given the somewhat weedy nature of both species, and increased rail and road traffic across the continent, it is quite possible either taxon could become naturalized elsewhere in the future. Some other indigenous Australian solanums have been recorded in self-sustaining populations from outside their usual ranges, e.g. *S. aviculare, S. capsiciforme, S. cinereum, S. laciniatum, S. oligacanthum, S. sturtianum* (Symon 1976, 1981, 1994 and pers. comm.) and *S. linearifolium* (B.J. Lepschi unpubl.). Notably, most taxa only persist at a locality for a few years without spreading far from the site of initial colonization, and then disappear. An exception is *S. cinereum* which has established naturalized populations.

Acknowledgements

I am grateful to David Symon for examining additional material in AD on my behalf, and to him, Greg Keighery and Terena Lally for their comments on the manuscript. I also thank the directors of AD for allowing me access to records from the ADHERB database.

References

Bitter, G. & Summerhayes, V.S. (1926). A new Solanum from Western Australia. Kew Bull. 1926: 116-119.

Black, J.M. (1926). "Flora of South Australia." Vol. 4. (Govt. Printer: Adelaide.)

Brown, R. (1810). "Prodromus florae Novae Hollandiae et Insulae van-Diemen." (J. Taylor: London.)

Conn, B.J. (1992). Solanum. In: Harden, G.J. (ed.). "Flora of New South Wales." Vol. 3. (New South Wales Univ. Press: Sydney.)

Grieve, B.J. & Blackall, W.E. (1975). "How To Know Western Australian Wildflowers." Part IV. (Univ. Western Australia Press: Perth.)

Lepschi, B.J. & Symon, D.E. (in press). A preliminary cladistic analysis of Australasian *Solanum* and *Lycianthes*. Solanaceae IV: Proceedings of the IV International Solanaceae Conference.

Meadly, G.R.W. (1960). Afghan Thistle. J. Dept. Agric. Western Australia 1: 401-402.

Parsons, W.T. & Cuthbertson, E.G. (1992). "Noxious Weeds of Australia." (Inkata Press: Melbourne.)

Paterson, J.G. (1967). Watch out for Afghan Thistic. J. Dept. Agric. Western Australia 8: 166-167.

Purdie, R.W., Symon, D.E. & Haegi, L.A.R. (1982). Solanaceae. In: George, A.S. (ed.). "Flora of Australia." Vol. 29. (Austral. Govt. Publ. Service: Canberra.)

Symon, D.E. (1976). The establishment and spread of Solanum cinereum R.Br., Narrawa Burr, in South Australia. S. Austral. Naturalist 51: 28-29.

Symon, D.E. (1981). A revision of the genus Solanum in Australia. J. Adelaide Bot. Gard. 4: 1-367.

Symon, D.E. (1986). Solanum. *In:* Jessop, J. & Toelken, H.R. (eds). "Flora of South Australia." Part 3. (South Austral. Govt. Printing Division: Adelaide.)

Symon, D.E. (1994). "Kangaroo Apples - Solanum Sect. Archaesolanum." (Privately published: Adelaide.)

B.J. Lepschi

Western Australian Herbarium, Department of Conservation and Land Management, PO Box 104, Como, Western Australia 6152.